

DELAYED EQUATION FOR CHARGED RIGID NONRELATIVISTIC BALL

Alexander A. Vlasov

High Energy and Quantum Theory
Department of Physics
Moscow State University
Moscow, 119899
Russia

Simple expression for self-force acting on radiating rigid charged ball is derived (Sommerfeld ball). It is shown that appropriate delayed equation of motion has solutions in general differ from that for Sommerfeld sphere - there are no "radiationless" solutions, but there are oscillating without damping solutions though self-force has nonzero value.

03.50.De

Long time ago to calculate back-reaction of electromagnetic field on moving with acceleration a charged classical (not quantum) particle of finite size Sommerfeld [1] considered two models - uniformly charged rigid sphere and uniformly charged rigid ball and found out explicit expressions for electromagnetic self-forces.

These expressions in nonrelativistic limit are much simpler then the original ones. Thus for uniformly charged rigid sphere (with radius a , total charge Q and mass m), moving along trajectory $\vec{R} = \vec{R}(t)$, the equation of motion with self-force, which include the back-reaction of electromagnetic radiation, is the known delayed equation (see, for ex., [2-5])

$$m\dot{\vec{v}}(t) = \vec{F}_{self} + \vec{F}_{ext},$$

$$\vec{F}_{self} = \eta [\vec{v}(t - 2a/c) - \vec{v}(t)] \quad (1)$$

here $\vec{v} = d\vec{R}/dt$, $\eta \equiv \frac{Q^2}{3ca^2}$, and $\vec{F}_{ext}$ - some external force.

This equation is often considered in literature (see, for ex., [2-5, 6, 7]) and has the following significant features:

1.) there are no "runaway" solutions - all solutions of (1) for zero external force exponentially tend to zero for $t \rightarrow \infty$ (are damped);

2.) there are "radiationless" solutions - solutions with zero value of the self-force, though the body moves with acceleration. In particular, for harmonic oscillations appropriate radiationless frequencies have discrete values.

As for charged rigid ball with radius a and total charge Q , till now there were not known the compact forms of nonrelativistic self-force and the following expansion in powers of a/cT was used:

$$\vec{F}_{self}(t) = -\frac{24Q^2}{ac^2} \cdot \sum_{n=0}^{\infty} \frac{(-2a)^n}{c^n n! (n+2)(n+3)(n+5)} \cdot \frac{d^{n+1}}{dt^{n+1}} \vec{v}(t) \quad (1)$$

One can derive this result as from the strict Sommerfeld result so from the well-known general Jackson form for a self-force in nonrelativistic, linear in velocity and its derivatives approximation for spherically symmetrical body [8]:

$$\vec{F}_{self}(t) = -\frac{2}{3c^2} \cdot \sum_{n=0}^{\infty} \frac{(-1)^n}{c^n n!} \cdot \gamma_{n-1} \cdot \frac{d^{n+1}}{dt^{n+1}} \vec{v}(t) \quad (2)$$

here γ_{n-1} - are the form-factors, do not depending from the time in approximation under consideration:

$$\gamma_{n-1} = \int d\vec{r} d\vec{r}' \rho \rho' |\vec{r} - \vec{r}'|^{n-1} \quad (3)$$

The density of the charge ρ for rigid uniformly charged ball is

$$\rho(\vec{r}) = \rho_0 \cdot \theta(a - r), \quad \rho_0 \equiv \frac{Q}{4\pi a^3/3} \quad (4)$$

With this density the form-factors γ_{n-1} in (2) are calculated as

$$\gamma_{n-1} = \int d\vec{r} d\vec{r}' \rho \rho' |\vec{r} - \vec{r}'|^{n-1} = \frac{36Q^2}{a} \cdot \frac{(2a)^n}{(n+2)(n+3)(n+5)} \quad (5)$$

Using (5) in (2), we get the result (1).

For our goal - to construct compact form of self-force in case under consideration - it will be more convenient to use another representation of Jackson result (2-3).

It is not too complicate to notice that the expansion (2-3) can be rewritten in the form of Taylor expansion of delayed acceleration $\vec{a} = \vec{\ddot{v}}$ of a body:

$$\vec{F}_{self}(t) = -\frac{2}{3c^2} \int \int d\vec{r} d\vec{r}' \frac{\rho \rho'}{|\vec{r} - \vec{r}'|} \sum_{n=0}^{\infty} \frac{1}{n!} \left(\frac{-|\vec{r} - \vec{r}'|}{c} \cdot \frac{d}{dt} \right)^n \vec{a}(t) =$$

$$= -\frac{2}{3c^2} \int \int d\vec{r} d\vec{r}' \frac{\rho \rho'}{|\vec{r} - \vec{r}'|} \vec{a}(t - \frac{|\vec{r} - \vec{r}'|}{c}) \quad (5)$$

After substitution of ρ in form of (4) into (5) and integration, the expression (5) reduces to

$$\begin{aligned} \vec{F}_{self}(t) = -k \left[\frac{2}{3} \left(\frac{a}{c} \right)^3 \vec{R}(t) - \left(\frac{a}{c} \right)^2 \left(\vec{R}^{[-1]}(t) + \vec{R}^{[-1]}(t - 2\frac{a}{c}) \right) - \right. \\ \left. - 2\frac{a}{c} \vec{R}^{[-2]}(t - 2\frac{a}{c}) + \vec{R}^{[-3]}(t) - \vec{R}^{[-3]}(t - 2\frac{a}{c}) \right] \end{aligned} \quad (6)$$

here

$$\vec{R}^{[-n]}(t) \equiv \int dt_1 \dots \int_{t_{n-1}}^{t_n} dt_n \vec{R}(t_n)$$

and

$$k \equiv \frac{3Q^2 c^3}{a^6}$$

The equation of motion of Sommerfeld ball then reads:

$$\ddot{\vec{R}}(t) = \vec{F}_{self}(t)/m + \vec{F}_{ext}(t)/m \quad (7)$$

To compare features of equation (7) with that for Sommerfeld sphere we consider the case of zero external force and search for solutions of (7) in the form of harmonic oscillations with frequency w , in general being complex number:

$$\vec{R}(t)^{[-3]} = \vec{A} \exp(iwt) \quad (8)$$

Consequently eq.(7) reduces to the transcendental equation which for new dimensionless variable $W \equiv iw\delta$ ($\delta \equiv 2a/c$) is

$$\frac{1}{k^*} W^5 = -\frac{1}{12} W^3 + \frac{1}{4} W^2 - 1 + \left(1 + \frac{1}{2} W \right)^2 \exp(-W) \quad (9)$$

here $k^* \equiv \frac{3Q^2 c^3}{ma^6} \cdot \frac{(2a)^5}{c^5} = \frac{96Q^2}{mc^2 a}$ - is dimensionless parameter.

R.H.S. of (9) is proportional to the value of the self-force (6), calculated for harmonic oscillations (8).

Consider the following cases.

1.

Let $Im(W) = 0$, i.e. the frequency w is purely imaginary: $w = iz$.

Then the function in R.H.S. of eq. (9) has zero value for $W = 0$ and for $W > 0$, ($z < 0$) its derivative with respect to W has negative sign:

$$\frac{d \text{ R.H.S.}}{dW} = W/2 \left(1 - W/2 - (1 + W/2) \exp(-W) \right) < 0$$

Thus this function is negative for $W > 0$. Consequently the equation (9), where L.H.S. is positive for $W > 0$, has no solutions for $W > 0$, ($z < 0$). Thus there are no solutions for $R(t)$ of the form

$$R \sim \exp(-zt), \quad z < 0$$

- i.e. there are no exponentially increasing with time t solutions.

So for Sommerfeld ball as for Sommerfeld sphere there are no exponentially "running away solutions".

2.

Let $\text{Re}(W) = 0$, $W \equiv iZ$, i.e. the frequency w is purely real number.

Then eq. (9) can be put in the form

$$iZ^5/k^* - iZ^3/12 + Z^2/4 + 1 = \exp(-iZ)(1 + iZ/2)^2 \quad (10)$$

Multiplication of (10) on its complex conjugate leads to the relation

$$(Z^5/k^* - Z^3/12)^2 = 0 \quad (11)$$

It has the solution

$$Z = \pm \sqrt{k^*/12} \quad (12)$$

Turning back to the eq.(10) with (12), we come to the result

$$Z^2/4 + 1 = \exp(-iZ)(1 + iZ/2)^2 \quad (13)$$

Reducing the R.H.S. of (13) to the form

$$(Z^2/4 + 1) \exp(-iZ + i2\Phi), \quad \text{tg}(\Phi) = Z/2$$

we get from (13)

$$Z = 2\Phi, \quad \text{or} \quad \text{tg}(Z/2) = Z/2 \quad (14)$$

So relations (14) and (12) taking together yield the solution of our problem:

if $Z = Z_n$, $n = 1, 2, 3, \dots$ - the discrete solutions of the transcendental eq. (14), then parameter k^* must satisfy the condition

$$\frac{k^*}{12} \equiv \frac{8Q^2}{mc^2a} = (Z_n)^2$$

Thus for these frequencies (and values of parameter k^*) there are harmonic oscillating solutions of ball equation (7), free of radiation damping.

This class of solutions is absent for Sommerfeld sphere.

On the other hand, taking

$$Z^5/k^* \rightarrow 0 \tag{15}$$

one can try to find whether there are the radiationless solutions (for which $\vec{F}_{self} = 0$ or $\left(R.H.S. \text{ of (9)}\right)_{W=iZ} = 0$) for the ball equation (7). Equation (11) due to (15) gives the immediate answer - there are no such solutions.

Thus one can see that Sommerfeld ball and Sommerfeld sphere models provide us with different behavior of moving charged body. For the Sommerfeld ball there are no "radiationless" solutions but there are solutions with not damped oscillations though the value of the self-force differs from zero.

REFERENCES

1. A.Sommerfeld, Gottingen Nachrichten, 29 (1904), 363 (1904), 201 (1905).
2. T.Erber, Fortschr.Phys., 9, 342 (1961).
3. P.Pearle, in *Electromagnetism*, ed. D.Tepliz, Plenum, NY, 1982, p.211.
4. S.Parrott, *Relativistic Electrodynamics and Differential Geometry*, Springer-Verlag, NY, 1987.
5. F.Rohrlich, Am.J.Physics, 65(11), 1051 (1997), Phys.Rev., D60, 084017 (1999)
6. Al. A. Vlasov, Vestnik Mosk. Univer., Fizika, N 5, 17 (1998), N 6, 15 (2001)
7. Alexander. A.Vlasov, in "Photon: old problems in light of new ideas", p. 126, ed. Valeri V. Dvoeglazov, Nova Sci. Publ., NY, 2000. E-print Archive: physics/9911059, 9912051, 0004026, 0103065, 0110034.
8. J. D. Jackson, *Classical Electrodynamics*, 3rd ed., Wiley, NY, 1999.